



# FOREST INSECT & DISEASE MANAGEMENT

USDA • FOREST SERVICE • NORTHERN REGION

State & Private Forestry • Missoula, MT 59801

79-2, *Revised*

3450

June 1980

## R-1 FOREST INSECT AND DISEASE DAMAGE SURVEY SYSTEM

By

Wayne E. Bousfield, Entomologist




### INTRODUCTION

The purpose of insect and disease damage surveys is to estimate losses caused by various pests, so that the land manager can prescribe appropriate management action.

Systematic ground data collection systems provide to the land manager loss data that can be used for cost benefit analysis, management plans, environmental impact statements, and for projecting losses at a forest level. Damage survey data could be integrated into timber management plans to fill voids in stand models to predict losses. This handbook describes field data collection procedures and data recording necessary to operate the supporting computer program called "INDIDS" (Insect and Disease Damage Survey).

### SURVEY TYPES

Usually, the land manager has only one specific pest in mind when conducting damage loss surveys. This system, however, provides for four basic pest types: (1) bark beetles, (2) defoliators, (3) disease, (4) miscellaneous insect and damage problems. Data can be collected for all four types at the same time if desired, but analyzed separately using the computer program "INDIDS" for the different options. The program provides detailed mensurational data of stands by species, size class, and damage class for each designated survey type and aggregates other problems in one column. Detailed statistics on other problems can be obtained by running the option desired. Each sample tree in the survey, however, is limited to three pest problems or severity.




For example, a tree could be attacked by bark beetles, have mistletoe, and be top killed by a defoliator. Another tree could be recorded as having moderate defoliation, be lightly top killed, and have root rot. This flexibility allows for several alternatives in data collection, and four options to summarize detailed statistics.

Periodic annual increment is also calculated by a growth model. If growth loss estimates caused by a defoliator are desired and the stand recently became infested, then radial growth measurements covering the period during the infestation and prior to the infestation are measured, and adjusted mean growth between host and nonhost trees is tested by covariance analysis. If differences are significant, expected periodic annual increment is computed using the growth model. Areas of chronic defoliation cannot be evaluated in this manner, however, if top kill is present, growth loss calculations are based on presence of top kill.

#### FIELD PROCEDURES


Areas examined must first be delineated into stands using standard photo interpretation (P.I.) guides explained in TM handbooks. Use existing stand delineations if the area is National Forest. Obtain either 2-inch or 2.6-inch/mile topographic maps of the area, and resource photography to transfer stand boundaries to field maps. Sample points should occur on a 5- by 10-chain grid to sample a stand satisfactorily. Stands less than 40 acres should be sampled with greater intensity depending on desired sampling error. The following diagram shows how plots could be placed in a 100-acre stand.



Because of the clumped distribution of bark beetle killed trees in a stand, sampling intensities should be greater for attacked trees than undamaged trees to obtain similar sampling errors. INDIDS provides for different sampling intensities.

When conducting bark beetle surveys, uninfested or undamaged tree information need only be taken on a sample of the total plots. However, at least 10 plots in each stand should be sampled for undamaged stand statistics. For example, if a total of 20 plots are planned for the survey then record data on all trees within the plot on the even-numbered plots, and on the odd-numbered plots record trees only if bark beetles have attacked or some other damaging agent is involved. Upon completion of the survey, record the number of plots where only unattacked trees were recorded (columns 67-68) and the total number of plots (columns 72-74). Diameters of all trees in the plot are recorded to the nearest one-tenth of an inch. Tree heights are measured to the nearest 1 foot on a sample basis. Twenty-five percent of all trees over 5 inches d.b.h. on each plot (figure 1) are measured for height. After the number of trees have been determined, select trees clockwise from north to be measured for height. A different sampling method may be incorporated to obtain height estimates.


Figure 1--Height Measurements of Sample Trees




No heights are measured for trees less than 5 inches d.b.h. Regression estimates based on diameters and heights of sample trees are used to obtain heights for unmeasured trees using a log transformation of the data. If periodic increment is desired, then habitat type must be obtained for each plot, and the last 10 years' radial growth measured for each tree that was sampled for height. Ten-year radial growth is measured to nearest one-tenth inch and interpolated to the nearest one-hundredth inch. If growth loss is desired in the case of defoliators, then X-year and Y-year growth measurements on increment cores explained later must be recorded. Douglas-fir, grand fir, subalpine fir, and spruce are considered the host species when infestations of spruce budworm or tussock moth occur. No provisions are provided for the user to designate different host species, although it could be done through computer programming. Ponderosa pine, white pine, limber or lodgepole pine will be considered nonhost species for covariance testing. Larch will not be considered, because it is both a host of budworm and larch casebearer which makes it difficult to determine years infested. If nonhost species are a minor part of the forest, then measure growth of all nonhost trees on the plot.

The supporting computer program "INDIDS" will handle both variable and fixed plot sampling including strip plots. Trees less than 5 inches d.b.h. are only sampled when they occur within a 1/300-acre or 6.8-foot radius fixed plot. A maximum of four best management trees under 5 inches d.b.h. will be recorded on the 300th-acre fixed plot. Excess trees are considered culls and will not be recorded. Trees greater than 5 inches d.b.h. are sampled from either variable plot method or fixed plot of one-tenth of an acre or larger plot. The following examples explain sample tree selection.

#### Variable plot





### RECORDING DATA FOR INDIDS

#### HEADER CARD

The following is an explanation of how to record sample tree data on the field data sheet (Exhibit 1).

Identification code: (1-3) Unique code which identifies header card with data cards, usually the Forest code followed by a number. For example, (Ident code 1 0 3) where 10 is the Forest code for Flathead and 3 for that particular stand. This number is used to identify cards from the same stand only.

Forest: (4-19) Alpha or numeric name of Forest, Park, Indian reservation, BLM, etc. Example F L A T H E A D

Stand or Area: (20-35) Use compartment, subcompartment, and stand number or drainage. Any combination of numeric or alpha characters that will identify the areas surveyed.

Examples 7 4 7 1 2 0 or R O C K C R E E K # 2

<u>Survey code</u>	(36-37)	<u>Survey type</u>
01	=	Bark beetle survey
02	=	Defoliator survey
03	=	Disease survey
04	=	Miscellaneous insect and damaging problems

Observers: (52-66) Names or initials of observers.

Type plot: (69)

1 = Variable plot for trees >5 inches d.b.h. In addition, trees <5 inches are recorded on the 1/300-acre plot.

2 = Fixed plot for trees >5 inches d.b.h. In addition, trees <5 inches d.b.h. are recorded on the 1/300-acre plot.

Size plot: (70-71) Enter BA factor or, 4 for  $\frac{1}{4}$ , 5 for  $\frac{1}{5}$  for fixed plots.  
Example of BAF 40 would be recorded as 4 0, or  
1/5-acre fixed plot would be recorded as 0 5.

Undamaged tree plots: (67-68) Enter number plots where undamaged tree data was recorded. If undamaged tree data was recorded on all plots then undamaged tree plots would equal total plots.

Total plots: (72-74) Enter total number of plots used in sampling the stand including undamaged stand plots. Include all plots even if some had no trees.

Date: (75-80) Enter month, day, and year.

#### Plot cards

Ident code: (1-3) Use same as header card. Example: 1 0 3

Plot: (5-6) Plot number in sequence. Do not duplicate plot numbers within a stand. Example 0 1 for first plot.

HAB: (8-10) Habitat type for plot. Use Montana valid habitat type numbers. Example: 5 2 1 for grand fir/Clintonia uniflora phase (appendix I).

Species: (12-14) Use valid species code. Left adjusted. Appendix II.  
Examples White pine W P  
Larch L

D.b.h.: (16-18) Enter d.b.h. to nearest one-tenth inch right adjusted. For example, a tree 12.3 inches in diameter is recorded as 1 2 3.

Height: The height of trees sampled for height measurements is entered to the nearest foot, right adjusted.

Age: (24-26) or tree number if identity is needed. Age measurements are optional (to be added later to Program).


10-year growth: (28-30) Last 10 years' radial growth in inches (two decimals assumed) right adjusted. Example of 1.05 inches 1 0 5

Example of .23 inches 2 3

Y-year growth: (32-34) Radial growth measurements for the period of infestation. Not less than 2 years and not to exceed the last 10 years (two decimals assumed, right adjusted) (figure 2).

X-year growth: (36-38) Radial growth measurement of the period of normal growth (two decimals assumed, right adjusted) (figure 3).

Figure 3



#### INSECT OR DISEASE CODES


Columns: (40-47)

Examine each tree and assign a damage code to each tree. Be sure to use the 0 code if no damaging agents are found. Use only valid codes

(appendix III). Definitions of the 0 code must be placed in first column (tree is undamaged). Do not place 0 in first column (40-41) and another code in second column (43-44). It makes no difference in order of type of codes used.

#### INDIDS Deck Set Up for 2 Stands

A maximum of 99 stands can be run at one time. At present each stand has a limit of 400 sample trees. Exhibit 2 shows examples of output from INDIDS.





For each option INDIDS computes, the trees per acre (TA), basal area (BA), cubic foot volume per acre (CFV), board foot volume (BFV), actual periodic annual increment in cubic feet per acre per year (APAI), and expected periodic annual increment (EPAI) for each species by damage class and four diameter classes. In addition, percent of stand damaged is computed in terms of trees per acre, basal area cubic foot volume, and board foot volume. Diameter of the tree of mean basal area is calculated by species and damage class.

## Appendix Montana forest habitat types

ANP code <sup>1</sup>	Abbreviation	Scientific names	Habitat types and phases	Common names
010	SCREE			
000		PINUS FLEXILIS CLIMAX SERIES		
040	PIFL/AGSP h.t.	Pinus flexilis/Agropyron spicatum h.t.	limber pine/bluebunch wheatgrass	
050	PIFL/FEID h.t.	Pinus flexilis/Festuca idahoensis h.t.	limber pine/Idaho fescue	
051	-FEID phase	-Festuca idahoensis phase	-Idaho fescue phase	
052	-FESC phase	-Festuca scabrella phase	-rough fescue phase	
070	PIFL/JUCQ h.t.	Pinus flexilis/Juniperus communis h.t.	limber pine/common juniper	
100		PINUS PONDEROSA CLIMAX SERIES		
110	PIPO/AND h.t. <sup>2</sup>	Pinus ponderosa/Andropogon spp. h.t.	ponderosa pine/bluestem	
130	PIPO/AGSP h.t.	Pinus ponderosa/Agropyron spicatum h.t.	ponderosa pine/bluebunch wheatgrass	
140	PIPO/FEID h.t.	Pinus ponderosa/Festuca idahoensis h.t.	ponderosa pine/Idaho fescue	
141	-FEID phase	-Festuca idahoensis phase	-Idaho fescue phase	
142	-FESC phase	-Festuca scabrella phase	-rough fescue phase	
160	PIPO/PUTR h.t.	Pinus ponderosa/Purshia tridentata h.t.	ponderosa pine/bitterbrush	
161	-AGSP phase	-Agropyron spicatum phase	-bluebunch wheatgrass phase	
162	-FEID phase	-Festuca idahoensis phase	-Idaho fescue phase	
170	PIPO/SYAL h.t.	Pinus ponderosa/Symphoricarpos albus h.t.	ponderosa pine/snowberry	
171	-SYAL phase	-Symphoricarpos albus phase	-snowberry phase	
172	-BERE phase	-Berberis repens phase	-creeping oregon grape phase	
180	PIPO/PRVI h.t.	Pinus ponderosa/Prunus virginiana h.t.	ponderosa pine/chokecherry	
181	-PRVI phase	-Prunus virginiana phase	-chokecherry phase	
182	-SHCA phase	-Shepherdia canadensis phase	-buffaloberry phase	
200		PSEUDOTSUGA MENZIESII CLIMAX SERIES		
210	PSME/AGSP h.t.	Pseudotsuga menziesii/Agropyron spicatum h.t.	Douglas-fir/bluebunch wheatgrass	
220	PSME/FEID h.t.	Pseudotsuga menziesii/Festuca idahoensis h.t.	Douglas-fir/Idaho fescue	
230	PSME/FESC h.t.	Pseudotsuga menziesii/Festuca scabrella h.t.	Douglas-fir/rough fescue	
250	PSME/VACA h.t.	Pseudotsuga menziesii/Vaccinium caespitosum h.t.	Douglas-fir/dwarf huckleberry	
260	PSME/PHMA h.t.	Pseudotsuga menziesii/Physocarpus malvaceus h.t.	Douglas-fir/ninebark	
261	-PHMA phase	-Physocarpus malvaceus phase	-ninebark phase	
262	-CARU phase	-Calamagrostis rubescens phase	-pinegrass phase	
280	PSME/VAGL h.t.	Pseudotsuga menziesii/Vaccinium globulare h.t.	Douglas-fir/blue huckleberry	
281	-VAGL phase	-Vaccinium globulare phase	-blue huckleberry phase	
282	-ARUV phase	-Arctostaphylos uva-ursi phase	-kinnikinnick phase	
283	-XETE phase	-Xerophyllum tenax phase	-beargrass phase	
290	PSME/LIBO h.t.	Pseudotsuga menziesii/Linnaea borealis h.t.	Douglas-fir/twinflower	
291	-SYAL phase	-Symphoricarpos albus phase	-snowberry phase	
292	-CARU phase	-Calamagrostis rubescens phase	-pinegrass phase	
293	-VAGL phase	-Vaccinium globulare phase	-blue huckleberry phase	
310	PSME/SYAL h.t.	Pseudotsuga menziesii/Symphoricarpos albus h.t.	Douglas-fir/snowberry	
311	-AGSP phase	-Agropyron spicatum phase	-bluebunch wheatgrass phase	
312	-CARU phase	-Calamagrostis rubescens phase	-pinegrass phase	
313	-SYAL phase	-Symphoricarpos albus phase	-snowberry phase	
320	PSME/CARU h.t.	Pseudotsuga menziesii/Calamagrostis rubescens h.t.	Douglas-fir/pinegrass	
321	-AGSP phase	-Agropyron spicatum phase	-bluebunch wheatgrass phase	
322	-ARUV phase	-Arctostaphylos uva-ursi phase	-kinnikinnick phase	
323	-CARU phase	-Calamagrostis rubescens phase	-pinegrass phase	
324	-PIPO phase	-Pinus ponderosa phase	-ponderosa pine phase	
330	PSME/CAGE h.t.	Pseudotsuga menziesii/Carex geyeri h.t.	Douglas-fir/elk sedge	
340	PSME/SPBE h.t.	Pseudotsuga menziesii/Spiraea betulifolia h.t.	Douglas-fir/white spiraea	
350	PSME/ARUV h.t.	Pseudotsuga menziesii/Arctostaphylos uva-ursi h.t.	Douglas-fir/kinnikinnick	
360	PSME/JUCQ h.t.	Pseudotsuga menziesii/Juniperus communis h.t.	Douglas-fir/common juniper	
370	PSME/ARCO h.t.	Pseudotsuga menziesii/Arnica cordifolia h.t.	Douglas-fir/heartleaf arnica	
380	PSME/SYOR h.t. <sup>2</sup>	Pseudotsuga menziesii/Symphoricarpos oreophilus h.t.	Douglas-fir/mountain snowberry	
400		PICEA CLIMAX SERIES		
410	PICEA/EQAR h.t.	Picea/Equisetum arvense h.t.	spruce/common horsetail	
420	PICEA/CLUN h.t.	Picea/Clintonia uniflora h.t.	spruce/queencup beadlily	
421	-VACA phase	-Vaccinium caespitosum phase	-dwarf huckleberry phase	
422	-CLUN phase	-Clintonia uniflora phase	-queencup beadlily phase	
430	PICEA/PHMA h.t.	Picea/Physocarpus malvaceus h.t.	spruce/ninebark	
440	PICEA/GATR h.t.	Picea/Galium triflorum h.t.	spruce/sweet-scented bedstraw	
450	PICEA/VACA h.t.	Picea/Vaccinium caespitosum h.t.	spruce/dwarf huckleberry	
460	PICEA/SEST h.t.	Picea/Senecio streptanthifolius h.t.	spruce/cleft-leaf groundsel	
461	-PSME phase	-Pseudotsuga menziesii phase	-Douglas-fir phase	
462	-PICEA phase	-Picea phase	-spruce phase	
470	PICEA/LIBO h.t.	Picea/Linnaea borealis h.t.	spruce/twinflower	
480	PICEA/SMST h.t.	Picea/Smilacina stellata h.t.	spruce/starry Solomon's seal	
500		ABIES GRANDIS CLIMAX SERIES		
510	ABGR/XETE h.t.	Abies grandis/Xerophyllum tenax h.t.	grand fir/beargrass	
520	ABGR/CLUN h.t.	Abies grandis/Clintonia uniflora h.t.	grand fir/queencup beadlily	
521	-CLUN phase	-Clintonia uniflora phase	-queencup beadlily phase	
522	-ARNU phase	Aralia nudicaulis phase	-wild sarsaparilla phase	
523	-XETE phase	-Xerophyllum tenax phase	-beargrass phase	
590	ABGR/LIBO h.t.	Abies grandis/Linnaea borealis h.t.	grand fir/twinflower	
591	-LIBO phase	-Linnaea borealis phase	-twinflower phase	
592	-XETE phase	-Xerophyllum tenax phase	-beargrass phase	

## Appendix 1 continued

11

ADP code <sup>1</sup>	Abbreviation	Scientific names	Habitat types and phases	Common names
THUJA Plicata CLIMAX SERIES				
501				
530	THPL/CLUN h.t.	Thuja plicata/Clintonia uniflora h.t.		western redcedar/queencup beadlily
531	-CLUN phase	-Clintonia uniflora phase		-queencup beadlily phase
532	-ARMJ phase	-Aralia nudicaulis phase		-wild sarsaparilla phase
533	-MEFE phase	-Menziesia ferruginea phase		-menziesia phase
550	THPI/CHHO h.t.	Thuja plicata/Opiopanax horridum h.t.		western redcedar/devil's club
TSUGA heterophylla CLIMAX SERIES				
570	TSHE/CLUN h.t.	Tsuga heterophylla/Clintonia uniflora h.t.		western hemlock/queencup beadlily
571	-CLUN phase	-Clintonia uniflora phase		-queencup beadlily phase
572	-ARMJ phase	-Aralia nudicaulis phase		-wild sarsaparilla phase
ABIES lasiocarpa CLIMAX SERIES				
600				
Lower subalpine h.t.s				
610	ABLA/OPHO h.t.	Abies lasiocarpa/Opiopanax horridum h.t.		subalpine fir/devil's club
620	ABLA/CLUN h.t.	Abies lasiocarpa/Clintonia uniflora h.t.		subalpine fir/queencup beadlily
621	-CLUN phase	-Clintonia uniflora phase		-queencup beadlily phase
622	-ARMJ phase	-Aralia nudicaulis phase		-wild sarsaparilla phase
623	-VACA phase	-Vaccinium caespitosum phase		-dwarf huckleberry phase
624	-XETE phase	-Xerophyllum tenax phase		-beargrass phase
625	-MEFE phase	-Menziesia ferruginea phase		-menziesia phase
630	ABLA/GATR h.t.	Abies lasiocarpa/Gallium triflorum h.t.		subalpine fir/sweet-scented bedstraw
640	ABLA/VACA h.t.	Abies lasiocarpa/Vaccinium caespitosum h.t.		subalpine fir/dwarf huckleberry
650	ABLA/CACA h.t.	Abies lasiocarpa/Calamagrostis canadensis h.t.		subalpine fir/bluejoint
651	-CACA phase	-Calamagrostis canadensis phase		-bluejoint phase
653	-GATR phase	-Gallium triflorum phase		-sweet-scented bedstraw phase
654	-VACA phase	-Vaccinium caespitosum phase		-dwarf huckleberry phase
660	ABLA/LIBO h.t.	Abies lasiocarpa/Linnaea borealis h.t.		subalpine fir/twinflower
661	-LIBO phase	-Linnaea borealis phase		-twinflower phase
662	-XETE phase	-Xerophyllum tenax phase		-beargrass phase
663	-VASC phase	-Vaccinium scoparium phase		-grouse whortleberry phase
670	ABLA/MEFE h.t.	Abies lasiocarpa/Menziesia ferruginea h.t.		subalpine fir/menziesia
680	TSME/MEFE h.t.	Tsuga mertensiana/Menziesia ferruginea h.t.		mountain hemlock/menziesia
690	ABLA/XETE h.t.	Abies lasiocarpa/Xerophyllum tenax h.t.		subalpine fir/beargrass
691	-VAGL phase	-Vaccinium globulare phase		-blue huckleberry phase
692	-VASC phase	-Vaccinium scoparium phase		-grouse whortleberry phase
710	TSME/XETE h.t.	Tsuga mertensiana/Xerophyllum tenax h.t.		mountain hemlock/beargrass
720	ABLA/VAGL h.t.	Abies lasiocarpa/Vaccinium globulare h.t.		subalpine fir/blue huckleberry
730	ABLA/VASC h.t.	Abies lasiocarpa/Vaccinium scoparium h.t.		subalpine fir/grouse whortleberry
731	-CARU phase	-Calamagrostis rubescens phase		-pinegrass phase
732	-VASC phase	-Vaccinium scoparium phase		-grouse whortleberry phase
733	-THOY phase	-Thalictrum occidentale phase		-western meadowrue phase
740	ABLA/ALSI h.t.	Abies lasiocarpa/Alnus sinuata h.t.		subalpine fir/Sitka alder
750	ABLA/CARU h.t.	Abies lasiocarpa/Calamagrostis rubescens h.t.		subalpine fir/pinegrass
770	ABLA/CLPS h.t.	Abies lasiocarpa/Clematis pseudopalmata h.t.		subalpine fir/virgin's bower
780	ABLA/ARCO h.t.	Abies lasiocarpa/Arnica cordifolia h.t.		subalpine fir/heartleaf arnica
790	ABLA/CAGE h.t. <sup>2</sup>	Abies lasiocarpa/Carex geyeri h.t.		subalpine fir/elk sedge
791	-CAGE phase	-Carex geyeri phase		-elk sedge phase
792	-PSME phase	-Pseudotsuga menziesii phase		-Douglas-fir phase
Upper subalpine h.t.s				
810	ABLA/RIMO h.t. <sup>2</sup>	Abies lasiocarpa/Ribes montigenum h.t.		subalpine fir/mountain gooseberry
820	ABLA-PIAL/VASC h.t.	Abies lasiocarpa-Pinus albicaulis/Vaccinium scoparium h.t.		subalpine fir-whitebark pine/grouse whortleberry
830	ABLA/LUHI h.t.	Abies lasiocarpa/Luzula hitchcockii h.t.		subalpine fir/smooth wood-rush
831	-VASC phase	-Vaccinium scoparium phase		-grouse whortleberry phase
832	-MEFE phase	-Menziesia ferruginea phase		-menziesia phase
840	TSME/LUHI h.t. <sup>2</sup>	Tsuga mertensiana/Luzula hitchcockii h.t.		mountain hemlock/smooth wood-rush
841	-VASC phase	-Vaccinium scoparium phase		-grouse whortleberry phase
842	-MEFE phase	Menziesia ferruginea phase		-menziesia phase
Timberline h.t.s				
890				
850	PIAL-ABLA h.t.s	Pinus albicaulis-Abies lasiocarpa h.t.s		whitebark pine-subalpine fir
860	LALY-ABLA h.t.s	Larix laricina-Abies lasiocarpa h.t.s		alpine larch-subalpine fir
870	PIAL h.t.s	Pinus albicaulis h.t.s		whitebark pine
PINUS contorta CLIMAX SERIES				
900				
910	PICO/PUTR h.t.	Pinus contorta/Purshia tridentata h.t.		lodgepole pine/bitterbrush
920	PICO/VACA c.t.	Pinus contorta/Vaccinium caespitosum c.t.		lodgepole pine/dwarf huckleberry
930	PICO/LIBO c.t.	Pinus contorta/Linnaea borealis c.t.		lodgepole pine/twinflower
940	PICO/VASC c.t.	Pinus contorta/Vaccinium scoparium c.t.		lodgepole pine/grouse whortleberry
950	PICO/CARU c.t.	Pinus contorta/Calamagrostis rubescens c.t.		lodgepole pine/pinegrass

Total Number of Habitat Types = 64

Total Number of Habitat Types, Phase, and Pinus contorta Community Type Categories = 105

<sup>1</sup> Automatic data processing codes for National Forest System use.<sup>2</sup> Minor type in Montana; described in other study areas.

APPENDIX IISpecies Codes

GF	=	Grand fir
AF or SAF	=	Subalpine fir
J	=	Juniper
L or WL	=	Larch
S	=	Spruce
WBL, WBP, WLP,	=	White bark or limber pine
LP or LPP	=	Lodgepole pine
WP	=	Western white pine
PP	=	Ponderosa pine
DF	=	Douglas-fir
Y	=	Yew
C or WRC	=	Western redcedar
WH or H	=	Western hemlock
MH	=	Mountain hemlock
B	=	Birch
ASH	=	Ash
CW	=	Cottonwood
ASP	=	Aspen
OH	=	Other hardwoods
AL	=	Alpine larch

APPENDIX IIIDamage Codes

Code

- 0 Healthy tree, no insect, disease, or damage problem
- 1 Unknown or natural mortality

Bark Beetles

- 2 Current beetle attack
- 3 Last year's attack
- 4 Older attack
- 5 Unsuccessful attack
- 6 Current strip attack
- 7 Older strip attack
- 8 Current secondary bark beetle attack
- 9 Older secondary bark beetle attack

Defoliation

- 11 Light defoliation
- 12 Moderate defoliation
- 13 Heavy defoliation
- 14 Light top kill
- 15 Moderate top kill
- 16 Heavy top kill
- 17 Defoliator caused mortality

Disease

- 20 Mistletoe infected
- 21 Mistletoe mortality
- 22 Root rot infected
- 23 Current root rot mortality
- 24 Older root rot mortality
- 25 Branch canker and stem rust
- 26 Stem canker and stem rust
- 27 Mortality - stem rust
- 28 Light needle disease infection
- 29 Moderate needle disease infection
- 30 Heavy needle disease infection
- 31 Butt and stem decay

## OTHER INSECT AND DAMAGING PROBLEMS

- 40 Shoot damaging insects
- 41 Stem boorers
- 42 Gall insects
- 43 Sucking insects (aphids and scales)
- 44 Webworms
- 45 Light winter damage
- 46 Moderate winter damage
- 47 Heavy winter damage
- 48 Animal damage
- 49 Spike tops
- 50 Other damaging agents

Ident. code	(1-3)			
Forest	(4-19)	Slope	Aspect	Elev
Stand or Area	(20-35)	Type plot	(69)	Physio Site
Survey code	(36-37)	Size plot	(70-71)	Hab
Type	(38-51)			
Observers	(52-66)	Soil		
Green tree plots	(67-68)	Total plots	(72-74)	Date
				(75-76) (77-78) (79-80)

[illegible]

FOREST INSECT AND DISEASE DAMAGE SURVEY

STAND  
FOREST

CONTINUATION SHEET

EXHIBIT I, continued

16

Ident. code (1-3)  
Plot HAB Specimen D.b.b. Height Age Growth 10-yr.  
(5-6) (8-10) (12-14) (16-18) (20-22) (24-26) (28-30) (32-34) (36-38) (40-42) (44-46) (48-50)  
disease codes Insect or



Exhibit 2. - BARK BEETLE SURVEY  
--SPECIES TOTAL--

3/20/80

RD

TEST 1

101

DIAMETER CLASS		UNDAM. CLP. STAND	CLP. ATTACK	LAST YEAR'S ATTACK	OLDER ATTACK	UNSUCC. ATTACK	CUR. STRIP ATTACK	OLDER STRIP ATTACK	CUR. SEC. ATTACK	OLDER SFC. ATTACK	OTHER I+D PROB.	OTHER CAM. AGENTS	OTHER MOFT.	STAND TOTAL
0- 4.9	TA	300.0	.0	.0	.0	.0	.0	.0	.0	.0	200.0	.0	.0	500.0
	BA	.0	.0	.0	.0	.0	.0	.0	.0	.0	25.1	.0	.0	25.1
5- 8.9	TA	168.6	.0	.0	.0	.0	.0	.0	.0	.0	116.5	.0	.0	275.1
	BA	60.0	.0	.0	.0	.0	.0	.0	.0	.0	26.7	.0	.0	86.7
	CFA	703.3	.0	.0	.0	.0	.0	.0	.0	.0	289.0	.0	.0	992.3
	BFA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
	APAI	64.9	.0	.0	.0	.0	.0	.0	.0	.0	21.1	.0	.0	76.0
	EPAI	64.9	.0	.0	.0	.0	.0	.0	.0	.0	25.6	.0	.0	80.5
9-11.9	TA	21.8	21.8	.0	.0	.0	.0	.0	.0	.0	85.1	.0	21.8	128.6
	BA	13.3	13.3	.0	.0	.0	.0	.0	.0	.0	53.3	.0	13.3	80.0
	CFA	374.1	357.0	.0	.0	.0	.0	.0	.0	.0	1466.0	.0	307.1	2147.2
	BFA	1743.7	1654.1	.0	.0	.0	.0	.0	.0	.0	6476.1	.0	1125.7	9345.5
	APAI	14.3	.0	.0	.0	.0	.0	.0	.0	.0	34.3	.0	.0	48.6
	EPAI	14.3	11.0	.0	.0	.0	.0	.0	.0	.0	45.3	.0	.0	59.6
12 +	TA	5.3	.0	15.4	.0	.0	.0	.0	.0	.0	38.5	9.3	15.4	76.6
	BA	13.3	.0	13.3	.0	.0	.0	.0	.0	.0	53.3	13.3	13.3	93.3
	CFA	409.5	.0	448.4	.0	.0	.0	.0	.0	.0	1783.7	409.5	398.3	3040.0
	BFA	2048.8	.0	2278.0	.0	.0	.0	.0	.0	.0	9094.1	2048.8	1872.7	15253.6
	APAI	6.1	.0	.0	.0	.0	.0	.0	.0	.0	21.8	4.0	.0	27.8
	EPAI	10.6	.0	11.0	.0	.0	.0	.0	.0	.0	34.0	7.1	.0	55.6
TOTALS	TA	489.7	21.8	15.4	.0	.0	.0	.0	.0	.0	440.0	9.3	37.2	982.2
	BA	86.7	13.3	13.3	.0	.0	.0	.0	.0	.0	158.5	13.3	26.7	265.1
	CFA	1486.9	357.0	448.4	.0	.0	.0	.0	.0	.0	3538.7	409.5	705.4	6179.5
	BFA	7792.4	1654.1	2278.0	.0	.0	.0	.0	.0	.0	15570.2	2048.8	2998.5	24599.2
	APAI	75.3	.0	.0	.0	.0	.0	.0	.0	.0	77.2	4.0	.0	152.5
	EPAI	79.9	11.0	11.0	.0	.0	.0	.0	.0	.0	104.9	7.1	.0	195.8

TA = Trees per acre  
BA = Basal area  
CFA = Cubic ft. volume per acre  
BFA = Board ft volume per acre

APAI = Actual periodic annual increment  
EPAI = Expected periodic annual increment  
(PAI) Cubic foot production per acre per year without damage

## Exhibit 2 continued

HARK REEF TIE SURVEY  
---SPECIES LP---

3/20/80

R0

TEST 1

100

DIAMETER	CLAS	UACAM.	CUP.	LAST	YEAR'S	OLDER	INSUC.	CUP.	OLDER	CUP.	OLDER	OTHER	OTHER	OTHER	STAND
CLAS	STAND	ATTACH	ATTACH	ATTACH	ATTACH	ATTACH	ATTACH	ATTACH	ATTACH	ATTACH	ATTACH	PROG.	DAM.	MORT.	TOTAL
0-4.0	TA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
	BA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
	TA	105.7	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	105.7
	BA	26.7	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	26.7
5-8.0	PFA	564.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	564.0
	MFA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
	APAT	40.7	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	40.7
	EPAT	40.7	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	40.7
	TA	21.8	21.8	.0	.0	.0	.0	.0	.0	.0	.0	21.8	.0	.0	43.5
	BA	13.3	13.3	.0	.0	.0	.0	.0	.0	.0	.0	13.3	.0	.0	26.7
9-11.9	PFA	374.1	357.9	.0	.0	.0	.0	.0	.0	.0	.0	357.9	.0	.0	731.9
	MFA	1655.1	1655.1	.0	.0	.0	.0	.0	.0	.0	.0	1655.1	.0	.0	3398.7
	APAT	14.3	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	14.3
	EPAT	14.3	11.0	.0	.0	.0	.0	.0	.0	.0	.0	11.0	.0	.0	25.3
	TA	.0	.0	15.4	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	15.4
	BA	.0	.0	13.3	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	13.3
12+	PFA	.0	.0	448.4	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	448.4
	MFA	.0	.0	2238.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	2238.0
	APAT	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
	EPAT	.0	.0	11.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	11.0
	TA	127.5	21.8	15.4	.0	.0	.0	.0	.0	.0	.0	21.8	.0	.0	164.6
	BA	30.0	13.3	13.3	.0	.0	.0	.0	.0	.0	.0	13.3	.0	.0	66.7
TOTALS	PFA	958.0	357.9	448.4	.0	.0	.0	.0	.0	.0	.0	357.9	.0	.0	1744.3
	MFA	1743.7	1655.1	2238.0	.0	.0	.0	.0	.0	.0	.0	1655.1	.0	.0	5636.8
	APAT	55.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	55.0
	EPAT	55.0	11.0	11.0	.0	.0	.0	.0	.0	.0	.0	11.0	.0	.0	77.0

Exhibit 2 continued

DIAMETER OF THE TREE OF MEAN BASAL AREA BY DAMAGE CLASS

3/20/80

PP - 100

YEAR 1

100

SPECIES	UNDA- M. STAND	CUP. ATTACK	LAST YEAR'S ATTACK	OLDER ATTACK	UN- SUC. ATTACK	CUR. STRIP ATTACK	OLDER STRIP ATTACK	CUR. SEC. ATTACK	OLDER SEC. ATTACK	OTHER I+D PROB.	OTHER DAM. AGENTS	OTHER MORT.	STAND TOTAL
RF	.0	.0	.0	.0	.0	.0	.0	.0	.0	19.2	.0	.0	19.2
AF	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
J	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
I	.0	.0	.0	.0	.0	.0	.0	.0	.0	9.8	.0	.0	9.8
E	.1	.0	.0	.0	.0	.0	.0	.0	.0	9.4	.0	.0	6.3
UCD	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
IP	7.6	10.6	12.6	.0	.0	.0	.0	.0	.0	10.6	.0	.0	6.6
GP	.0	.0	.0	.0	.0	.0	.0	.0	.0	11.9	.0	12.6	12.2
DP	6.8	.0	.0	.0	.0	.0	.0	.0	.0	4.8	.0	.0	5.6
DF	3.4	.0	.0	.0	.0	.0	.0	.0	.0	7.2	16.2	10.6	5.9
Y	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
WH	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
WH	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
R	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
ASH	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
CW	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
SP	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
OH	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
AL	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
TOTAL	5.0	10.6	12.6	.0	.0	.0	.0	.0	.0	8.1	16.2	11.5	7.0

CROWN COMPETITION FACTOR = 322.9861

## Exhibit 2 continued

BARK BEETLE SURVEY  
--SPECIES TOTAL--

3/20/80

		FD		TEST 1		100									
DIAMETER		UNIAM.	CUR.	LAST	YEAR'S	OLDER	UNsuc.	CUR.	OLDER	CUR.	OLDER	OTHER	OTHER	OTHER	STAND
CLASS		STAND	ATTACK	ATTACK	ATTACK	ATTACK	ATTACK	ATTACK	ATTACK	ATTACK	ATTACK	PROB.	DAM.	MORT.	TOTAL
0-2.9	TA	300.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	300.0
	BA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
	CFA														
3-4.9	TA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	200.0	.0	.0	200.0
	BA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	25.1	.0	.0	25.1
	CFA														
5-6.9	TA	125.5	.0	.0	.0	.0	.0	.0	.0	.0	.0	116.5	.0	.0	242.0
	BA	26.7	.0	.0	.0	.0	.0	.0	.0	.0	.0	26.7	.0	.0	53.3
	CFA	346.1	.0	.0	.0	.0	.0	.0	.0	.0	.0	289.0	.0	.0	645.1
7-8.9	TA	33.1	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	33.1
	BA	13.3	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	13.3
	CFA	347.2	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	347.2
9-10.9	TA	21.8	21.8	.0	.0	.0	.0	.0	.0	.0	.0	67.8	.0	21.8	111.3
	BA	13.3	13.3	.0	.0	.0	.0	.0	.0	.0	.0	40.0	.0	13.3	66.7
	CFA	374.1	357.9	.0	.0	.0	.0	.0	.0	.0	.0	1016.9	.0	307.1	1696.1
11-12.9	TA	.0	.0	15.4	.0	.0	.0	.0	.0	.0	.0	17.3	.0	15.4	48.1
	BA	.0	.0	13.3	.0	.0	.0	.0	.0	.0	.0	13.3	.0	13.3	40.0
	CFA	.0	.0	448.4	.0	.0	.0	.0	.0	.0	.0	451.1	.0	398.3	1297.6
13-14.9	TA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	13.2	.0	.0	13.2
	BA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	13.3	.0	.0	13.3
	CFA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	388.7	.0	.0	388.7
15-16.9	TA	9.3	.0	.0	.0	.0	.0	.0	.0	.0	.0	18.6	9.3	.0	27.9
	BA	13.3	.0	.0	.0	.0	.0	.0	.0	.0	.0	26.7	13.3	.0	40.0
	CFA	679.5	.0	.0	.0	.0	.0	.0	.0	.0	.0	890.1	409.5	.0	1299.6
17-18.9	TA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
	BA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
	CFA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
19+	TA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	6.6	.0	.0	6.6
	BA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	13.3	.0	.0	13.3
	CFA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	504.9	.0	.0	504.9
TOTALS	TA	489.7	21.8	15.4	.0	.0	.0	.0	.0	.0	.0	440.0	9.3	37.2	982.2
	BA	46.7	13.3	13.3	.0	.0	.0	.0	.0	.0	.0	158.5	13.3	26.7	265.1
	CFA	1466.9	357.9	448.4	.0	.0	.0	.0	.0	.0	.0	3538.7	409.5	705.4	6179.5

## Exhibit 2 continued

DEFOLIATOR SURVEY  
--SPECIES TOTAL--

3/20/80

		NO		TEST 1		100									
DIAMETER CLASS		UNDAM. STAND	LIGHT OFF.	MOD. DEF.	HEAVY DEF.	LIGHT TOP KILL	MOD. TOP KILL	HEAVY TOP KILL	DEF. MORT.	OTHER I+D PROR.	OTHER MORT.	OTHER DAM. AGENTS	STAND TOTAL		
0- 4.9	TA	300.0	.0	.0	100.0	.0	.0	.0	.0	100.0	.0	.0	500.0		
	BA	.0	.0	.0	12.6	.0	.0	.0	.0	12.6	.0	.0	25.1		
5- 8.9	TA	158.6	.0	.0	.0	63.6	.0	.0	.0	116.5	.0	.0	275.1		
	BA	40.0	.0	.0	.0	13.3	.0	.0	.0	26.7	.0	.0	66.7		
	CFA	703.3	.0	.0	.0	111.4	.0	.0	.0	289.0	.0	.0	992.3		
	BFA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0		
	APAI	54.9	.0	.0	.0	6.0	.0	.0	.0	21.1	.0	.0	76.0		
	EPAI	54.9	.0	.0	.0	10.5	.0	.0	.0	25.6	.0	.0	80.5		
9-11.9	TA	21.8	.0	20.6	.0	.0	.0	.0	.0	85.1	.0	21.8	128.6		
	BA	13.3	.0	13.3	.0	.0	.0	.0	.0	53.3	.0	13.3	80.0		
	CFA	374.1	.0	376.3	.0	.0	.0	.0	.0	1466.0	.0	307.1	2147.2		
	BFA	1743.7	.0	1723.3	.0	.0	.0	.0	.0	6476.1	.0	1125.7	9345.5		
	APAI	14.3	.0	11.0	.0	.0	.0	.0	.0	34.3	.0	.0	48.6		
	EPAI	14.3	.0	11.0	.0	.0	.0	.0	.0	45.3	.0	.0	59.6		
12 +	TA	9.3	.0	.0	9.3	22.5	.0	6.6	.0	44.6	9.3	15.4	78.6		
	BA	13.3	.0	.0	13.3	26.7	.0	13.3	.0	53.3	13.3	13.3	93.3		
	CFA	409.5	.0	.0	409.5	798.3	.0	504.9	.0	1822.6	409.5	398.3	3040.0		
	BFA	2048.8	.0	.0	2048.8	3871.7	.0	2776.6	.0	9283.4	2048.8	1672.7	15253.6		
	APAI	6.1	.0	.0	4.0	9.7	.0	4.3	.0	17.7	4.0	.0	27.8		
	EPAI	10.6	.0	.0	7.1	17.0	.0	9.2	.0	37.9	7.1	.0	55.6		
TOTALS	TA	489.7	.0	20.6	109.3	86.1	.0	6.6	.0	340.1	9.3	37.2	962.2		
	BA	66.7	.0	13.3	25.9	40.0	.0	13.3	.0	145.9	13.3	26.7	265.1		
	CFA	1486.9	.0	376.3	409.5	909.6	.0	504.9	.0	3577.6	409.5	705.4	6179.5		
	BFA	3792.4	.0	1723.3	2048.8	3871.7	.0	2776.6	.0	15759.5	2048.8	2990.5	24599.2		
	APAI	75.3	.0	11.0	4.0	15.7	.0	4.3	.0	73.1	4.0	.0	152.5		
	EPAI	79.9	.0	11.0	7.1	27.5	.0	9.2	.0	108.8	7.1	.0	195.8		

## Exhibit 2 continued

DISEASE SURVEY  
SPECIES TOTAL

3/20/80

		RD		TEST 1		100												
DIAMETER CLASS		LACAM. STAND	MIS. INF.	MIS. MORT.	ROOT ROT INF.	CUP. ROOT ROT MORT.	OLDER ROOT ROT MORT.	BRANCH CANKER	STEM CANKER	STEM RUST MORT.	LIGHT NEEDLE DIS.	MOD. NEEDLE DIS.	HEAVY NEEDLE DIS.	BUTT, STEM DECAY	OTHER I+D PROB.	OTHER DAM. AGENTS	OTHER MORT.	STAND TOTAL
0-4.9	TA	300.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	200.0	.0	.0	500.0
	BA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	25.1	.0	.0	25.1
5-8.9	TA	158.6	63.6	.0	.0	.0	.0	.0	.0	.0	63.6	.0	.0	.0	116.5	.0	.0	275.1
	BA	40.0	13.3	.0	.0	.0	.0	.0	.0	.0	13.3	.0	.0	.0	26.7	.0	.0	66.7
	CFA	703.3	111.4	.0	.0	.0	.0	.0	.0	.0	111.4	.0	.0	.0	289.0	.0	.0	992.3
	RFA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0
	APAI	54.9	6.0	.0	.0	.0	.0	.0	.0	.0	6.0	.0	.0	.0	21.1	.0	.0	76.0
	EPAI	54.9	10.5	.0	.0	.0	.0	.0	.0	.0	10.5	.0	.0	.0	25.6	.0	.0	60.5
9-11.9	TA	21.8	21.8	.0	.0	.0	.0	39.0	17.3	.0	.0	25.5	.0	.0	42.3	.0	21.8	128.6
	BA	13.3	13.3	.0	.0	.0	.0	26.7	13.3	.0	.0	13.3	.0	.0	26.7	.0	13.3	80.0
	CFA	374.1	357.9	.0	.0	.0	.0	808.9	451.1	.0	.0	280.8	.0	.0	734.1	.0	307.1	2147.2
	RFA	1743.7	1655.1	.0	.0	.0	.0	3780.6	2125.6	.0	.0	972.2	.0	.0	3378.4	.0	1125.7	9345.5
	APAI	14.3	.0	.0	.0	.0	.0	12.6	12.6	.0	.0	10.7	.0	.0	11.0	.0	.0	48.6
	EPAI	14.3	11.0	.0	.0	.0	.0	23.6	12.6	.0	.0	10.7	.0	.0	22.0	.0	.0	59.6
12 +	TA	9.3	13.2	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	6.6	53.9	9.3	15.4	78.6
	BA	13.3	13.3	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	13.3	66.7	13.3	13.3	93.3
	CFA	409.5	388.7	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	504.9	2232.1	409.5	398.3	3040.0
	RFA	2048.8	1823.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	2776.6	11332.2	2048.8	1872.7	15253.6
	APAI	6.1	5.7	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	4.3	21.8	4.0	.0	27.8
	EPAI	10.6	10.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	9.2	45.0	7.1	.0	55.6
TOTALS	TA	489.7	98.6	.0	.0	.0	.0	39.0	17.3	.0	63.6	25.5	.0	6.6	412.7	9.3	37.2	982.2
	BA	66.7	40.0	.0	.0	.0	.0	26.7	13.3	.0	13.3	13.3	.0	13.3	145.1	13.3	26.7	265.1
	CFA	1486.9	957.9	.0	.0	.0	.0	808.9	451.1	.0	111.4	280.8	.0	504.9	3255.3	409.5	705.4	6179.5
	RFA	3752.4	3478.0	.0	.0	.0	.0	3780.6	2125.6	.0	.0	972.2	.0	2776.6	114710.5	2048.8	2998.5	4599.2
	APAI	75.3	11.7	.0	.0	.0	.0	12.6	12.6	.0	6.0	10.7	.0	4.3	53.9	4.0	.0	152.5
	EPAI	79.9	31.5	.0	.0	.0	.0	23.6	12.6	.0	10.5	10.7	.0	9.2	92.6	7.1	.0	195.8

Exhibit 2 continued

OTHER INSECT AND DAMAGING PROBLEM SURVEY  
-----SPECIES TOTAL-----

3/20/80

		RC		TEST 1		100											
DAMETER	CLASS	UNDAM.	SHOOT	STEM	GALL	SCALES.	WFR-	LIGHT	MOD.	HEAVY	ANIMAL	SPIKE	OTHER	OTHER	OTHER	STAND	
		STAND	INSECT	BORER	INSECT	APHID	WORMS	WIN.	WIN.	WIN.	DAM.	DAM.	PROB.	UAP.	MURT.	TOTAL	
1- 4.9	TA	300.0	100.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	100.0	.0	.0	500.0	
	BA	.0	12.6	.0	.0	.0	.0	.0	.0	.0	.0	.0	12.6	.0	.0	25.1	
1- 8.9	TA	150.6	.0	.0	52.9	.0	.0	.0	.0	.0	.0	.0	63.6	.0	.0	275.1	
	BA	40.0	.0	.0	13.3	.0	.0	.0	.0	.0	.0	.0	13.3	.0	.0	66.7	
	CFA	703.3	.0	.0	177.7	.0	.0	.0	.0	.0	.0	.0	111.4	.0	.0	992.3	
	RFA	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	.0	
	APAI	54.9	.0	.0	15.1	.0	.0	.0	.0	.0	.0	.0	6.0	.0	.0	76.0	
	EPAI	54.9	.0	.0	15.1	.0	.0	.0	.0	.0	.0	.0	10.5	.0	.0	80.5	
1-11.9	TA	21.8	.0	.0	20.6	.0	.0	.0	.0	.0	.0	.0	85.1	.0	21.8	128.6	
	BA	13.3	.0	.0	13.3	.0	.0	.0	.0	.0	.0	.0	53.3	.0	13.3	80.0	
	CFA	374.1	.0	.0	376.3	.0	.0	.0	.0	.0	.0	.0	1466.0	.0	307.1	2147.2	
	RFA	1743.7	.0	.0	1723.3	.0	.0	.0	.0	.0	.0	.0	6476.1	.0	1125.7	9345.5	
	APAI	14.3	.0	.0	11.0	.0	.0	.0	.0	.0	.0	.0	34.3	.0	.0	48.6	
	EPAI	14.3	.0	.0	11.0	.0	.0	.0	.0	.0	.0	.0	45.3	.0	.0	59.6	
2 +	TA	9.3	.0	.0	.0	.0	.0	.0	.0	.0	.0	9.3	44.6	9.3	15.4	78.6	
	BA	13.3	.0	.0	.0	.0	.0	.0	.0	.0	.0	13.3	53.3	13.3	13.3	93.3	
	CFA	409.5	.0	.0	.0	.0	.0	.0	.0	.0	.0	480.6	1751.6	409.5	398.3	3040.0	
	RFA	2146.6	.0	.0	.0	.0	.0	.0	.0	.0	.0	2445.7	8886.4	2048.8	1872.7	15253.6	
	APAI	6.1	.0	.0	.0	.0	.0	.0	.0	.0	.0	7.7	14.0	4.0	.0	27.8	
	EPAI	10.6	.0	.0	.0	.0	.0	.0	.0	.0	.0	7.7	37.3	7.1	.0	55.6	
TAI S	TA	409.7	100.0	.0	77.4	.0	.0	.0	.0	.0	.0	9.3	293.2	9.3	37.2	982.2	
	BA	66.7	12.6	.0	26.7	.0	.0	.0	.0	.0	.0	13.3	132.6	13.3	26.7	265.1	
	CFA	1486.9	.0	.0	553.9	.0	.0	.0	.0	.0	.0	480.6	3328.9	409.5	705.4	6179.5	
	RFA	7702.4	.0	.0	1723.3	.0	.0	.0	.0	.0	.0	2445.7	15362.5	2048.8	2998.5	24599.2	
	APAI	75.3	.0	.0	26.1	.0	.0	.0	.0	.0	.0	7.7	54.3	4.0	.0	152.5	
	EPAI	79.9	.0	.0	26.1	.0	.0	.0	.0	.0	.0	7.7	93.1	7.1	.0	155.8	